arXiv:1006.0173v1 [math.CO] 1 Jun 2010

Circular Sequences and the Diameter of Multipermutohedra

Sarang Aravamuthan*

June 2, 2010

Abstract

We derive bounds on the number of switches at an arbitrary set of positions in a circular sequence of permutations and relate them to the diameter of Multipermutohedra.

Keywords: Circular sequences, Multipermutohedra
MSC (2010): 05A05, 52B12

1 Introduction

A sequence of $\bar{N}+1$ (with $\bar{N}=\binom{N}{2}$) permutations, $\mathcal{P}=\langle \pi_0,\ldots,\pi_{\bar{N}}\rangle$, in the symmetric group $\mathcal{S}_N$ is called a *circular sequence* if $\pi_0 = 1\,2\ldots n$ is the identity permutation, $\pi_{\bar{N}} = n\,(n-1)\ldots 1$ is the reverse permutation and π_{i+1} differs from π_i by an adjacent transposition, i.e., $\pi_{i+1}=\pi_i\,(j,j+1)$ for some j such that $\pi_i(j)<\pi_i(j+1)$. These sequences are used for bounding the number of k-sets of a point configuration $\mathcal{X}$ in the plane (a k-set of $\mathcal{X}$ is a subset of size k separated by a line from the other points in $\mathcal{X}$). The connection between k-sets and circular sequences was first established by Goodman and Pollack [5]. For a subset $K\subseteq[(n-1)/2]$, one can also bound the number of k-sets, $k\in K$ over all point configurations $\mathcal{X}$. Such bounds were established for specific choices of K using this connection (see [1, 6]).

For the sequence $\mathcal{P}$, defined above, the process of moving from π_i to π_{i+1} is referred to as a *switch* $(\pi_i(j),\pi_i(j+1))$ at position j (note that the numbers being swapped are $\pi_i(j)$ and $\pi_i(j+1)$). We define $s_j(\mathcal{P})$ to be the total number of switches at position j. We count the total number of switches at a given set of positions $y=\{y_1,\ldots,y_n\}\subseteq[N-1]$, with $y_1<y_2<\cdots<y_n$ which is

$$s(\mathcal{P},y)=s_{y_1}(\mathcal{P})+\cdots+s_{y_n}(\mathcal{P}).$$

---

*Ignite R&D Labs, Tata Consultancy Services, Chennai, India.
sarang.aravamuthan@tcs.com

Our goal is to derive a bound on $s(\mathcal{P}, y)$ and show how it relates to the diameter of *multipermutohedra* (which, as we will see later, are essentially "permutohedra on multisets").

In the next section, we estimate $s(\mathcal{P}, y)$. In Section 3, we introduce the multipermutohedron and derive bounds on its diameter. In Section 4, we show how the diameter of multipermutohedra relates to a variant of the k-set problem.

We fix some notations. $[n]$ represents the set $\{1, 2, \ldots, n\}$. If x is not an integer, we write $[x]$ instead of $[\lfloor x \rfloor]$. For $S \subseteq [n]$ and $x \in \mathbb{R}^n$ define $x(S) := \sum\limits_{i \in S} x_i$ and denote the size of S by $|S|$.

A *composition* of n is a sequence of positive integers $\lambda := \langle \lambda_1, \ldots, \lambda_k \rangle$ with $\sum \lambda_i = n$.

Permutations in $\mathcal{S}_n$ will be represented as words. We call $e := 1\,2 \ldots N \in \mathcal{S}_N$ the identity permutation and $\bar{e} := N\,N-1 \ldots 1$ the reverse permutation.

2 A Bound on Circular Sequences

With $\mathcal{P}$ and y as defined above, we evaluate $s(\mathcal{P}, y)$ in the following way. Each $i \in [N]$ starts at position i in the identity permutation e and reaches position $N - i + 1$ in $\bar{e}$. Following the notation of [6], we refer to the positions $y_1, \ldots, y_n$ as *gates* and define c_i to be the number of gates that i must cross in order to reach position $N - i + 1$. c_i is interpreted as the cost in moving i to position $N - i + 1$. As $c_i = c_{N-i+1}$, the total cost $c(N)$ is

$$c(N) := \sum_{i=1}^{N} c_i = 2 \sum_{i=1}^{\lfloor \frac{N}{2} \rfloor} c_i.$$

Next, we define the vector $x = (x_1, \ldots, x_n)$ where x_j represents the distance of y_j from one of the ends, i.e.

$$x_j := \min\{y_j, N - y_j\}. \tag{1}$$

Let $(p_1, \ldots, p_n)$ be the permutation of $(x_1, \ldots, x_n)$ with $N/2 \geq p_1 \geq \cdots \geq p_n > 0$. Any number j such that $p_{i+1} < j \leq p_i$ has cost $c_j = i$ (we assume $p_{n+1} = 0$). Hence

$$c(N) = 2 \sum_{i=1}^{n} i(p_i - p_{i+1}) = 2p([n]) = 2x([n]).$$

We now amortize the cost $c(N)$ over $\mathcal{P}$. Each switch (i, j) across a gate is interpreted as contributing $+1$ or -1 to c_i according to whether i moves towards or away from position $N-i+1$. A switch (i,j) across a gate is called *good* if it contributes 1 to both c_i and c_j and *bad* if it contributes 1 to c_i (or c_j) and -1 to the other. We observe that every switch across a gate is either good or bad, i.e., must contribute $+1$ to the cost of at least one of the numbers being moved. This follows from the fact that $i<j$, so if i is in position $> N-i > N-j$, then j moves towards $N-j+1$.

Hence, good switches contribute 2 to $c(N)$ while bad switches contribute 0. Since $c(N) = 2x([n])$, the number of good switches is $x([n])$. Thus if $s_b(\mathcal{P}, y)$ is the number of bad switches, then

$$s(\mathcal{P}, y) = x([n]) + s_b(\mathcal{P}, y). \tag{2}$$

Thus, it suffices to estimate $s_b(\mathcal{P}, y)$, in order to bound $s(\mathcal{P}, y)$. Let l be the number of y_i smaller than $N/2$, i.e., $y_l < N/2$ while $y_{l+1} \geq N/2$, and let $r := n-l$. In other words, l and r represent the number of gates to the $l$eft and $r$ight of $N/2$ respectively. The following result bounds $s(\mathcal{P}, y)$.

Theorem 1 *Let*

$$s(y) = \min\{s(\mathcal{P}, y) \mid \mathcal{P} \text{ is a circular sequence}\}$$

be the minimum of the number of switches at a set of positions given by y over all circular sequences of permutations in $\mathcal{S}_N$. Then,

$$x([n]) \leq s(y) \leq x([n]) + lr.$$

Proof. From (2), we observe that $s(\mathcal{P}, y) \geq x([n])$ for any circular sequence $\mathcal{P}$. This proves the lower bound for $s(y)$.

To show the upper bound, we construct a circular sequence $\mathcal{P}$ such that the number of bad switches $s_b(\mathcal{P}, y) \leq lr$. We construct the sequence in two phases.

In the first phase, we move the numbers $1, \ldots, r$ to positions $y_r, y_{r-1}, \ldots y_1$ and the numbers $N, N-1, \ldots, N-l+1$ to positions $y_{r+1}, \ldots, y_n$ in the following way. Starting from e, we move 1 to position y_r through a sequence of switches $(1,i), i \leq y_r$. Next, 2 is moved to position y_{r-1} with the switches $(2,i), i \leq y_{r-1}$. Continuing this way, the first r and the last l numbers (in the order $N, N-1, \ldots, N-l+1$) are moved to positions $y_r, y_{r-1}, \ldots, y_1$ and $y_{r+1}, \ldots, y_n$ respectively.

The resulting permutation is

$$r+1 \ldots r| \ldots r-1| \ldots\ldots 2| \ldots 1| \ldots N| \ldots N-1| \ldots\ldots N-l+1| \ldots N-l$$

where the $|$ indicates the gate positions $y_1, \ldots, y_n$ and the numbers $r+1, \ldots, N-l$ are in the remaining positions in increasing order.

The second phase consists of moving $N-i+1$ to position i for $i = 1, 2, \ldots, \lfloor \frac{N}{2} \rfloor$. This is done by starting with N and moving it to the first position by switching it in succession with the numbers $1, \ldots, r$. This brings $1, \ldots, r$ to positions $y_{r+1}, y_r, \ldots, y_2$ and $r+1$ to position y_1 if $y_1 > 1$. Next we move $N-1$ to the second position. In general, suppose γ is a permutation in this sequence with the last i numbers in the first i positions and the first j numbers in the last j positions, and $\gamma(i+1) = k$, i.e.

$$\gamma = N\, N-1 \ldots N-i+1 \; k \; \ldots \; k-1| \ldots k-2| \ldots \ldots j+1| \ldots N-i \; j\, j-1 \ldots 1.$$

Here the numbers $j+1, \ldots, k-1$ are at the gate positions and the numbers $k, k+1, \ldots, N-i-1$ are in the remaining positions in increasing order. Switching $N-i$ in succession with $j+1, j+2, \ldots, k$ moves $N-i$ to position $i+1$, and $j+1$ to position $N-j$ resulting in the permutation

$$N\, N-1 \ldots N-i \; k+1 \ldots k| \ldots k-1| \ldots \ldots j+2| \ldots N-i-1 \; j+1\, j \ldots 1.$$

Repeating this operation, we finally obtain the permutation $\bar{e}$.

We observe that all switches in the second phase are good; for a switch (i, j) across a gate, the numbers $1, \ldots, (i-1)$ are already to the right of i. So i is at position $< N-i+1$. The same argument holds for j.

In the first phase, suppose $l \leq r$. Then the number of bad switches are at most $l(r-l)$ for moving the numbers $1, \ldots, r-l$ to positions $y_r, y_{r-1}, \ldots, y_{l+1}$; $(l-1) + (l-2) + \cdots + 0 = l(l-1)/2$ for moving the numbers $r-l+1, \ldots, r$ to positions $y_l, y_{l-1}, \ldots, y_1$ and $l + (l-1) + \cdots + 1 = l(l+1)/2$ for moving the numbers $N, N-1, \ldots, N-l+1$ to positions $y_{r+1}, \ldots, y_n$ adding up to a total of at most lr bad switches. The case $l > r$ is handled similarly. This proves the result. □

Example 2 Let $N = 8$ and $y = \{1, 4, 6, 7\}$. By (1), $x = (1, 4, 2, 1)$. Since $y_1 < N/2$ and $y_2 \geq N/2$, $l = 1$ and $r = n - l = 3$. By Theorem 1,

$$8 = \sum x_i \leq s(y) \leq 8 + lr = 11$$

To construct a circular sequence $\mathcal{P}$ with $s(\mathcal{P}, y) = 11$, we start with the identity permutation and move the numbers $1, 2, 8$ to positions y_3, y_2, y_4 leading to 3 4 5 2 6 1 8 7. Next we move 8 to position 1 by switching it in succession with the numbers $1, 2, 3$ giving 8 4 5 3 6 2 1 7. Next 7 is moved to the second place. We summarize by showing some of the permutations in this sequence.

$$\begin{array}{lclcl}
1\,|2\,3\,4\,|5\,6\,|7\,|8 & \xrightarrow{4} & \underline{3}\,|4\,5\,\underline{2}\,|6\,\underline{1}\,|\underline{8}\,|7 & \xrightarrow{3} & \\
8\,|\underline{4}\,5\,\underline{3}\,|6\,\underline{2}\,|\underline{1}\,|\underline{7} & \xrightarrow{3} & 8\,|7\,\underline{5\,4}\,|\underline{6}\,3\,|2\,|1 & \xrightarrow{1} & 8\,7\,6\,5\,4\,3\,2\,1.
\end{array}$$

where the numbers above the arrows indicate the number of switches required to move to the next permutation, the '|' show the positions where the switches are counted and the numbers being switched are underlined. □

We see that the lower bound for $s(y)$ is attained when all the gates are to the left (or right) of the middle. In this case, $r = 0$ (resp. $l = 0$) and the two bounds for $s(y)$ coincide. The upper bound is attained when $y = [N-1]$. In this case, $s(y) = \binom{N}{2}$, i.e. we count switches at all positions.

3 Circular Sequences and Multipermutohedra

The multipermutohedron is a generalization of the permutohedron P_N which is the convex hull of all permutations of the point $(1, 2, \ldots, N) \in \mathbb{R}^N$. A multipermutohedron is defined by taking the convex hull of all permutations of a *multiset*, that is, a "set" with repeated elements.

Let $b_1 < b_2 < \cdots < b_n$ be n distinct numbers ($n > 1$) and consider the multiset with k_i copies of b_i for $i = 1, \ldots, n$, i.e.

$$M := \{b_1^{k_1}, b_2^{k_2}, \ldots, b_n^{k_n}\} = \{a_1, \ldots, a_N\} \qquad k_i > 0,\ i = 1, \ldots, n \tag{3}$$

with $a_1 \le a_2 \le \cdots \le a_N$ and $N = \sum_{i=1}^{n} k_i$. Let $P(M) \subset \mathbb{R}^N$ be the polytope formed by taking the convex hull of all permutations of the point $(a_1, \ldots, a_N) \in \mathbb{R}^N$. We call $P(M)$ the *multipermutohedron* on the multiset M.

Multipermutohedra were first studied by Schoute [7], who constructed them all from the simplex by a sequence of simple operations of *expansion* and *contraction*. Independent treatments of multipermutohedra also appear in [2, 8]. The proofs for the assertions that follow can be found in [2].

The inequality description of $P(M)$ is a direct generalization of the description for the permutohedron.

$$P(M) = \{x \in \mathbb{R}^N \mid x([N]) = \sum_{i=1}^{N} a_i,\ x(S) \ge \sum_{i=1}^{|S|} a_i,\ \text{for all } S \subset [N]\ \}.$$

Thus if $M = \{1, 2, 2, 3\}$, then $P(M) \subset \mathbb{R}^4$ is given by

$$\begin{array}{rcll} x_1 + x_2 + x_3 + x_4 & = & 8 & \\ 1 \le x_i & \le & 3 & i = 1, \ldots, 4 \\ x_i + x_j & \ge & 3 & 1 \le i < j \le 4. \end{array}$$

Figure 1 shows $P(\{1, 2, 2, 3\})$. It has 14 facets corresponding to these 14 inequalities, which correspond, in turn, to the 14 ordered partitions of the set $\{1, 2, 3, 4\}$ into two parts.

Faces of $P(M)$ are products of lower dimensional multipermutohedra and correspond to ordered partitions of $[N]$. As in the permutohedron, the face lattice of the multipermutohedron does not depend on the numbers being permuted but only on their multiplicities. Henceforth we assume that, in (3), $b_i = i$



Figure 1: A 3-dimensional multipermutohedron $P(\{1,2,2,3\})$.

and rewrite M as

$$M = \{1^{k_1}, \dots, n^{k_n}\}. \tag{4}$$

Given $\sigma \in \mathcal{S}_N$, the vertex a_σ has 1's in positions $\sigma(1), \dots, \sigma(k_1)$, 2's in positions $\sigma(k_1+1), \dots, \sigma(k_1+k_2)$ and so on, i.e.

$$a_\sigma := (a_{\sigma^{-1}(1)}, \dots, a_{\sigma^{-1}(N)}). \tag{5}$$

Note that this correspondence is not one-to-one; several permutations will correspond to the same vertex of $P(M)$.

Adjacency of vertices on $P(M)$ is obtained by switching two components whose values are consecutive. The relation to permutations is as follows. For $j = 1, \dots, n$, define

$$y_j := k_1 + \cdots + k_j \tag{6}$$

and let $y_0 = 0$. We define *blocks* of consecutive integers $S_1, \dots, S_n$ by

$$S_j := [y_j] \setminus [y_{j-1}].$$

Given $\sigma, \pi \in \mathcal{S}_N$, the vertices a_σ and a_π are adjacent if $a_\pi = a_{\sigma(i,j)}$ for some transposition (i,j) with i and j in successive blocks. In other words, π is obtained from σ by switching $\sigma(i)$ and $\sigma(j)$ for some i and j in successive blocks and possibly permuting the values of $\sigma(i,j)$ within each block.

Example 3 Let $M = \{1,2,2,3\}$ as earlier. Let $e = 1\,2\,3\,4$ be the identity permutation. This corresponds to the vertex $a_e = (1,2,2,3)$ of $P(M)$. Multiplying e by transpositions (i,j) with i,j in successive blocks leads to the permutations $2\,1\,3\,4$, $3\,2\,1\,4$, $1\,4\,3\,2$ and $1\,2\,4\,3$. These in turn correspond to the vertices $(2,1,2,3)$, $(2,2,1,3)$, $(1,3,2,2)$ and $(1,2,3,2)$ that are adjacent to $(1,2,2,3)$ as shown in Figure 1. $\square$

For $\sigma, \pi \in \mathcal{S}_N$, we denote by $d(\sigma, \pi)$, the shortest distance between the vertices a_σ and a_π in $P(M)$. For convenience, we also denote the vertex a_σ in $P(M)$ by σ.

The *diameter* of $P(M)$ (denoted $\operatorname{diam}(P(M))$) is the largest of the distances between pairs of vertices of $P(M)$.

Proposition 4 *The vertex farthest from $\sigma \in P(M)$ is $\bar{\sigma} = \sigma(N)\,\sigma(N-1)\ldots\sigma(1)$. The diameter of $P(M)$ is $d(e, \bar{e})$.*

Proof. We first show that the vertex farthest from e in $P(M)$ is $\bar{e}$. This will follow if we can show that for any permutation π with $a_\pi \neq a_{\bar{e}}$, switching a pair of numbers $\pi(i) < \pi(j)$ with $i < j$ in successive blocks yields a permutation $\pi' = \pi(i,j)$ with $d(e, \pi') \geq d(e, \pi)$.

Suppose $d(e, \pi) > d(e, \pi')$. Let $i' = \pi(i)$ and $j' = \pi(j)$. Since π and π' are adjacent vertices in $P(M)$, $d(e, \pi) = d(e, \pi') + 1$. Consider the shortest path from e to π through π'. Let γ, α_1 be successive vertices on this path with $\alpha_1^{-1}(j') < \alpha_1^{-1}(i')$ in successive blocks while $\gamma^{-1}(i')$ and $\gamma^{-1}(j')$ are either in the same block or in successive blocks. Hence, in going from γ to α_1, we have moved j' to the left of i'. Let the vertices on the path from γ be $\gamma, \alpha_1, \ldots, \alpha_{p-1} = \pi', \alpha_p = \pi$. We construct a shorter path from γ to π, namely $\gamma, \beta_1, \ldots, \beta_{p-1} = \pi$ where for each m, $\beta_m = (i', j')\alpha_m$. We see that for each $m > 0$, β_{m+1} differs from β_m by a transposition across adjacent blocks showing that they are adjacent. Since β_1 switches i' and j' in α_1, we see that either $\beta_1 = \gamma$ or β_1 and γ are adjacent (this case subsumes the possibility of $\beta_1 = \alpha_1$). In any case we see that this path is shorter than our chosen path by at least 1. This contradicts our hypothesis that $d(e, \pi) > d(e, \pi')$ proving our claim.

Given a path from e to π in $P(M)$, multiplying the permutations in this path by σ yields a path from σ to $\sigma\pi$ of the same length. Hence the vertex farthest from σ is $\sigma\bar{e} = \bar{\sigma}$ and $d(\sigma, \bar{\sigma}) = d(e, \bar{e})$ proving the result. $\square$

We now relate the diameter of $P(M)$ to circular sequences. Let the multiset M and $y = (y_1, \ldots, y_{n-1})$ be given by (4) and (6) respectively. From the criterion for adjacency of vertices on $P(M)$, we observe that each circular sequence $\mathcal{P}$ corresponds to a path in $P(M)$ from e to $\bar{e}$ of length $s(\mathcal{P}, y)$. From Proposition 4, it follows that

$$\operatorname{diam}(P(M)) = \min\{s(\mathcal{P}, y) \mid \mathcal{P} \text{ is a circular sequence}\}.$$

Theorem 1 automatically translates to bounds on diameter of $P(M)$.

Corollary 5 *Let M be defined by (4) with $k_i > 0$ and let $y = (y_1, \ldots, y_{n-1})$ and $x = (x_1, \ldots, x_{n-1})$ be specified by (6) and (1). Then the diameter of $P(M)$ is bounded by*

$$x([n-1]) \leq \operatorname{diam}(P(M)) \leq x([n-1]) + lr$$

with l and r as defined in Theorem 1. □

Example 6 Consider the multiset $M = \{1, 2^3, 3^2, 4, 5\}$. The vectors y and x given by (6) and (1) are $y = (1, 4, 6, 7)$ and $x = (1, 4, 2, 1)$. Since $y_1 < N/2$ and $y_2 \geq N/2$, $l = 1$ and $r = n - 1 - l = 3$. By Corollary 5,

$$8 = \sum x_i \leq \mathrm{diam}\,(P(M)) \leq 8 + lr = 11.$$

A circular sequence $\mathcal{P}$ with $s(\mathcal{P}, y) = 11$ was constructed in Example 2 and this translates to a path of length 11 between e and $\bar{e}$ in $P(M)$. □

It's easy to see that the upper bound in Corollary 5 is attained for the permutohedron P_N which has a diameter of $\binom{N}{2}$. Also, for small n, it's possible to derive an explicit expression for diam$(P(M))$.

Proposition 7 *Let the multiset M be given by the composition $\langle k_1, \ldots, k_n \rangle$ of N. If $n = 2$ then $\mathrm{diam}\,(P(M)) = \min\{k_1, k_2\}$. When $n = 3$,*

$$\mathrm{diam}\,(P(M)) = \begin{cases} x_1 + x_2 & \text{if } k_1 \neq k_3; \\ x_1 + x_2 + 1 = 2k_1 + 1 & \text{if } k_1 = k_3. \end{cases}$$

Proof. For $n = 2$, the lower and upper bounds for the diameter in Corollary 5 are the same and the result follows. If $n = 3$ and $k_1 \neq k_3$ then it is easy to show by a careful choice of switches that the lower bound for the diameter is attainable. If $k_1 = k_3$, then by Corollary 5, the diameter is either $2k_1$ or $2k_1 + 1$. The lower bound is not attained because, for any circular sequence, the switches at positions k_1 and $N - k_1$ cannot all involve a number less than k_1 and a number greater than $N - k_1$, i.e., there must be at least one bad switch. This proves the result. □

We observe that since the diameter of $P(M)$ is obtained by counting the number of switches at certain positions in a circular sequence in $\mathcal{S}_N$, its value is at most the total number of switches i.e. $\binom{N}{2}$. It would be interesting to determine, in some form, all the integers in the set $[\binom{N}{2}]$ that are the diameters of multipermutohedra $P(M) \subset \mathbb{R}^N$ for some M given by (4).

4 Multipermutohedra and k-sets

We now relate the diameter of the multipermutohedron to arrangements of points on a plane. Let $\mathcal{X} \subset \mathbb{R}^2$ be a configuration of N points in general position on a plane, i.e., no three points of $\mathcal{X}$ lie on a line. For $k \leq \lfloor N/2 \rfloor$, we define a *left* (resp. *right*) *k-set* to be a set of k points of $\mathcal{X}$ that lie on the left (resp. right) of a line (with respect to a directed reference line, say the X-axis).

For a line parallel to the X-axis, we take the left of the line to be the open half-space above the line. A k-set of $\mathcal{X}$ is either a left k-set or a right k-set. Let $f_l(k, \mathcal{X})$, $f_r(k, \mathcal{X})$ and $f(k, \mathcal{X})$ denote the number of left k-sets, right k-sets and k-sets of $\mathcal{X}$ respectively.

For subsets $L, R \subseteq [(N-1)/2]$, we count the number of sets that appear as a left k-set for $k \in L$ or a right k-set for $k \in R$. We define

$$f(L, R, \mathcal{X}) := \sum_{k \in L \cap R} f(k, \mathcal{X}) + \sum_{k \in L \setminus R} f_l(k, \mathcal{X}) + \sum_{k \in R \setminus L} f_r(k, \mathcal{X}). \tag{7}$$

When $L = R \subseteq [(N-1)/2]$, we are counting the number of k-sets for $k \in L$ and we write $f(K, \mathcal{X})$ for $f(K, K, \mathcal{X})$. Our objective is to derive bounds for $f(L, R, \mathcal{X})$ in terms of the diameters of certain multipermutohedra.

Since the minimum (resp. maximum) of $f(L, R, \mathcal{X})$ is not affected by a slight perturbation of the points of $\mathcal{X}$, we assume that no two points of $\mathcal{X}$ lie on a line parallel to the X-axis. Then, for each $k \leq (N-1)/2$, there are two k-sets that are both left k- and right k-sets. Hence

$$f_l(k, \mathcal{X}) + f_r(k, \mathcal{X}) = f(k, \mathcal{X}) + 2. \tag{8}$$

Following the approach of Goodman and Pollack [5], we associate with $\mathcal{X}$ a circular sequence of permutations in the following way. Project the points of $\mathcal{X}$ onto a directed line ℓ parallel to the Y-axis. Label the projected points on ℓ in the order $1, \ldots, N$ and rotate ℓ counter clockwise. The projection of the points on ℓ determines a permutation $\pi \in \mathcal{S}_N$. The first k points $\pi(1), \ldots, \pi(k)$ form a left k-set while the last k points $\pi(N-k+1), \ldots, \pi(N)$ form a right k-set. When ℓ becomes perpendicular to a line joining a pair of points of $\mathcal{X}$, the order of the projection of these two points on ℓ is reversed. The new permutation differs from π by an adjacent transposition. When ℓ has rotated through an angle of 180^o, the order of the projected points becomes $N, N-1, \ldots, 1$. Thus $\mathcal{X}$ describes a circular sequence of permutations $\mathcal{P}(\mathcal{X}) = \langle e, \pi_1, \ldots, \pi_{\bar{N}} = \bar{e} \rangle$. If s_k is the number of switches at position k in $\mathcal{P}(\mathcal{X})$, then the number of left k-sets $f_l(k, \mathcal{X}) = s_k + 1$ and the number of right k-sets $f_r(k, \mathcal{X}) = s_{N-k} + 1$. Hence by (8), the number of k-sets $f(k, \mathcal{X}) = s_k + s_{N-k}$. By (7),

$$f(L, R, \mathcal{X}) = \sum_{k \in L} s_k + \sum_{k \in R} s_{N-k} + |L \cup R| - |L \cap R|. \tag{9}$$

The subsets L and R define a composition $\langle L, R \rangle = \langle k_1, \ldots, k_n \rangle$ of N where $n = |L| + |R| + 1$ and the elements of L are the partial sums $k_1 + k_2 + \cdots + k_i$ that are at most $(N-1)/2$ while the elements of R are the partial sums $k_n + k_{n-1} + \cdots + k_j$ that are at most $(N-1)/2$. Let $P(\langle L, R \rangle)$ be the multipermutohedron defined by this composition. Then the sequence $\mathcal{P}(\mathcal{X})$ describes a path in $P(\langle L, R \rangle)$ from e to $\bar{e}$ of length $\sum\limits_{k \in L} s_k + \sum\limits_{k \in R} s_{N-k}$ which must be at least its diameter. This gives a lower bound for $f(L, R, \mathcal{X})$.

To bound $f(L, R, \mathcal{X})$ from above, we consider the subsets $\bar{L} := [N/2] \setminus L$ and $\bar{R} := [(N-1)/2] \setminus R$. As before, these define a composition $\langle \bar{L}, \bar{R} \rangle$ of N and a multipermutohedron $P(\langle \bar{L}, \bar{R} \rangle)$. Since $\sum_{k=1}^{N-1} s_k = \binom{N}{2}$, we can rewrite (9) as

$$f(L, R, \mathcal{X}) = \binom{N}{2} - \left(\sum_{k \in \bar{L}} s_k + \sum_{k \in \bar{R}} s_{N-k} \right) + |L \cup R| - |L \cap R|.$$

This gives the upper bound for $f(L, R, \mathcal{X})$ and ties the diameter of $P(M)$ to k-sets in the following way.

Theorem 8 *Let $\mathcal{X}$ be a configuration of N points in $\mathbb{R}^2$ in general position and let $L, R \subseteq [(N-1)/2]$. Then the function $f(L, R, \mathcal{X}) - |L \cup R| + |L \cap R|$ is bounded from above and below by the diameter of multipermutohedra $P(\langle L, R \rangle)$ and $P(\langle \bar{L}, \bar{R} \rangle)$, that is,*

$$\operatorname{diam}(P(\langle L, R \rangle)) \leq f(L, R, \mathcal{X}) - |L \cup R| + |L \cap R| \leq \binom{N}{2} - \operatorname{diam}(P(\langle \bar{L}, \bar{R} \rangle)).$$

In particular,

$$\operatorname{diam}(P(\langle K, K \rangle)) \leq f(K, \mathcal{X}) \leq \binom{N}{2} - \operatorname{diam}(P(\langle \bar{K}, \bar{K} \rangle))$$

thus bounding the number of k-sets for $k \in K \subseteq [(N-1)/2]$ by diameters of multipermutohedra. □

The problem of k-sets and its relation to circular sequences has been extensively studied (see [1, 4, 5, 6]). In [6], k-sets are used to derive a lower bound on the number of convex quadrilaterals in a set of n points in the plane.

A lower bound for the number of k-sets of a point configuration in general position follows from Proposition 7 and the above theorem. As observed in [6, Example 8], this lower bound of $2k+1$ can actually be achieved by a configuration of N points ($N \geq 2k+1$) which consist of a regular $(2k+1)$-gon with the remaining points situated close to the center of the gon. Finding the upper bound for the number of k-sets seems to be a harder problem. Some estimates for this bound are given in [4, 3].

Acknowledgment. The author would like to thank L. J. Billera for helpful discussions.

References

[1] N. Alon and E. Györi, The number of small semispaces of a finite set of points in the plane, *J. Combinatorial Theory Ser. A* **41** (1986), 154-157.

`http://www.math.tau.ac.il/~nogaa/PDFS/Publications/The%20number%20of%20small%20semispaces%20of%20a%20finite%20set%20of%20points%20in%20the%20plane.pdf`

[2] L. J. Billera and A. Sarangarajan, The combinatorics of permutation polytopes, in L. Billera, C. Greene, R. Simion and R. Stanley, eds., *Algebraic Combinatorics, DIMACS Series in Discrete Mathematics and Theoretical Computer Science*, American Mathematical Society, 1994. `http://igm.univ-mlv.fr/~fpsac/FPSAC94/ARTICLES/billera.ps.gz`

[3] T. Dey, Improved bounds on planar k-sets and related problems, *Discrete Comput. Geom.*, **19** (1998), 373-382. `http://www.cse.ohio-state.edu/~tamaldey/paper/kset/paper.ps.gz`

[4] P. Erdös, L. Lovász, A. Simmons and E.G. Straus, Dissection graphs of planar point sets, in *A Survey of Combinatorial Theory*, (J.N. Srivastava et al., Eds.) 139-149, North Holland, Amsterdam 1973. `http://www.cs.elte.hu/~lovasz/scans/DissectGraph.pdf`

[5] J. E. Goodman and R. Pollack, On the combinatorial classification of nondegenerate configurations in the plane, *J. Combinatorial Theory Ser. A* **29** (1980), 220-235.

[6] L. Lovász, K. Vesztergombi, U. Wagner, E. Welzl, Convex Quadrilaterals and k-sets, in: J. Pach (Ed.), Towards a Theory of Geometric Graphs, *AMS Contemporary Mathematics* **342** (2004) 139-148. `http://www.cs.elte.hu/~lovasz/quadrilaterals.pdf`

[7] P. H. Schoute, Analytic treatment of the polytopes regularly derived from the regular polytopes, *Verhandelingen der Koninklijke Akademie van Wetenschappen te Amsterdam* **11**, No. 3, Johannes Müller, Amsterdam 1911.

[8] A. S. Schulz, *Polyedrische Charakterisierung von Scheduling Problemen*, Diploma Thesis, Technische Universität Berlin, Berlin, Germany, 1993.